

Implications of Yukawa Textures in the decay $H^+ \rightarrow W^+\gamma$ within the 2HDM-III

J. E. Barradas-Guevara,^{1,*} F. Cázarez-Bush,^{2,†} A. Cordero-Cid,^{3,‡} O.

Félix-Beltrán,^{3,§} J. Hernández-Sánchez,^{3,4,¶} and R. Noriega-Papaqui^{5,4,**}

¹*Fac. de Cs. Físico-Matemáticas, BUAP. Apdo. Postal 1364, C.P. 72000 Puebla, Pue., México.*

²*Instituto de Matemáticas, Universidad Nacional Autónoma de México,*

Cd. Universitaria, México, D.F. 01000, México.

³*Facultad de Ciencias de la Electrónica, Benemérita Universidad Autónoma de Puebla,*

Apdo. Postal 542, C. P. 72570 Puebla, Pue., México.

⁴*Dual C-P Institute of High Energy Physics, Puebla, Pue., México.*

⁵*Centro de Investigación en Matemáticas,*

Universidad Autónoma del Estado de Hidalgo,

Carr. Pachuca-Tulancingo Km. 4.5, C.P. 42184, Pachuca, Hgo., México.

(Dated: February 26, 2010)

We discuss the implications of assuming a four-zero Yukawa texture for the charged Higgs decay $H^+ \rightarrow W^+\gamma$ at one-loop level, within the context of the general 2-Higgs Doublet Model of Type III. We begin by presenting a detailed analysis of the charged Higgs boson couplings with heavy quarks and the resulting effects on its decays. In particular, we present the possible enhancement of the decay $H^+ \rightarrow W^+\gamma$, whose branching ratio could be of the order $\sim 10^{-1}$, for charged Higgs mass of the order 180 GeV. These parameters can still avoid the $B \rightarrow X_s\gamma$ constraint and the perturbativity bound. The production of charged Higgs bosons is also sensitive to the modifications of its couplings, and we evaluate the events rates at the LHC, including the ‘direct’ $c\bar{b} \rightarrow H^+ + c. c.$ and ‘indirect’ $q\bar{q}, gg \rightarrow t\bar{b}H^+ + c. c.$ production channels. We get an event rate of order 60 (120) for ‘direct’ (‘indirect’) production reaction, and including decay $H^+ \rightarrow W^+\gamma$ in the final state, with an integrated luminosity of $10^5 pb^{-1}$.

PACS numbers: 12.60.Cn, 12.60.Fr, 11.30.Er

*Electronic address: barradas@fcfm.buap.mx

†Electronic address: federico@matem.unam.mx

‡Electronic address: acordero@ece.buap.mx

§Electronic address: olga'f'lix@@ece.buap.mx

¶Electronic address: jaimeh@ece.buap.mx

**Electronic address: rnoriega@uaeh.edu.mx

I. INTRODUCTION

Detecting a charged Higgs boson during the imminent Large Hadron Collider (LHC) experimental running would constitute a clear evidence of physics beyond the Standard Model (SM) [1]. Charged Higgs bosons appear in many well motivated extensions of the SM, whose phenomenology has been widely studied over the years [2–4]. In particular, 2-Higgs Doublet Models (2HDMs), in both Supersymmetry (SUSY) and non-SUSY versions [5, 6], can be considered as a prototype of a Higgs sector that includes a charged Higgs boson ($H^\pm$). It is expected that the LHC will allow us to test the mechanism of Electro-Weak Symmetry Breaking (EWSB) and, in particular, to probe the properties of charged Higgs bosons, which represent a unique probe of a weakly-interacting theory, as is the case of the Minimal Supersymmetric Standard Model (MSSM) [5] and general 2HDMs of Type I, II, III and IV (2HDM-I, 2HDM-II, 2HDM-III and 2HDM-IV) [7], or whether strongly-interacting scenarios are instead realized, like in the old Technicolor models or the ones discussed more recently [8]. Ultimately, while many analyses in this direction can be carried out at the LHC, it will be a future International Linear Collider (ILC) [9] or Compact Linear Collider (CLIC) [10] which will have the definite word about exactly which mechanism of mass generation and which realization of it occurs in Nature.

The 2HDM-II has been quite attractive to date, in part because it coincides with the Higgs sector of the MSSM, wherein each Higgs doublet couples to the u - or d -type fermions separately¹. However, this is only valid at tree-level [12]. When radiative effects are included, it turns out that the MSSM Higgs sector corresponds to the most general version of the 2HDM, namely the 2HDM-III, whereby both Higgs fields couple to both quarks and leptons. Thus, we can consider the 2HDM-III as a generic description of physics at a higher scale (of order TeV or maybe even higher), whose low energy imprints are reflected in the Yukawa coupling structure. With this idea in mind, a detailed study of the 2HDM-III Yukawa Lagrangian was presented in Refs.[13, 14], under the assumption of a specific texture pattern [15], which generalizes the original model of Ref. [16]. Phenomenological implications of this model for the neutral Higgs sector, including Lepton Flavour Violation (LFV) and/or Flavour Changing Neutral Currents (FCNCs) have been presented in a previous work [17]. The extension of such an approach to investigate charged Higgs boson phenomenology was conducted in Ref. [14], which discussed the implications of this Yukawa texture for the charged Higgs boson properties (masses and couplings) and the resulting pattern

¹ Notice that there exist significant differences between the 2HDM-II and MSSM though, when it comes to their mass/coupling configurations and possible Higgs signals [11].

of charged Higgs boson decays and main production reactions at the LHC.

Decays of charged Higgs bosons have been studied in the literature, including the radiative modes $W^\pm\gamma, W^\pm Z^0$ [18, 19], mostly within the context of the 2HDM-II or its SUSY incarnation (i.e., the MSSM), but also by using an effective Lagrangian extension of the 2HDM [20]. More recently, within an extension of the MSSM with one Complex Higgs Triplet (MSSM+1CHT) [21, 22]. While the decay mode $H^+ \rightarrow W^+\gamma$ is forbidden at the tree level due to electromagnetic gauge invariance, the $H^+ \rightarrow W^+Z$ decay can be induced at this order in models including Higgs triplets or more complicated representations [2, 21]. In spite of their suppressed branching ratios, these decay modes are very interesting, due to the fact that their experimental study may provide important information concerning the underlying structure of the gauge and scalar sectors; these channels have a clear signature and might be at the reach of current and future particle colliders. Charged Higgs boson production at hadron colliders was studied long ago [23, 24] and, more recently, systematic calculations of production processes at the LHC have been presented [25].

Current bounds on the mass of a charged Higgs boson have been obtained at Tevatron, by studying the top decay $t \rightarrow b H^+$, which already eliminates large regions of the parameter space [26], whereas LEP2 bounds imply that, approximately, $m_{H^+} > 80$ GeV [27, 28], rather model independently. Concerning theoretical limits, tree-level unitarity bounds on the 2HDM Higgs masses have been studied in generic 2HDMs and in particular an upper limit for the charged Higgs mass of 800 GeV or so can be obtained, according to the results of Ref. [29].

In this paper we extend previous studies, to include the decay $H^+ \rightarrow W^+\gamma$. This paper is organized as follows. In section II, we discuss the Higgs-Yukawa sector of the 2HDM-III, in particular, we derive the expressions for the charged Higgs boson couplings to heavy fermions. Then, in section III, we review the expressions for the decays $H^+ \rightarrow W^+\gamma$ at one-loop level and numerical results are presented for some 2HDM-III scenarios, defined for phenomenological purposes. Actual LHC event rates for the main production mechanisms at the LHC are given in section IV. These include the s -channel production of charged Higgs bosons through $c\bar{b}(\bar{c}b)$ -fusion [30] and the multi-body more $q\bar{q}, gg \rightarrow t\bar{b}H^- + \text{c.c.}$ (charge conjugated). These mechanisms depend crucially on the parameters of the underlying model and large deviations could be expected in the 2HDM-III with respect to the 2HDM-II. Finally, we summarize our results and present the conclusions in section V.

II. THE CHARGED HIGGS BOSON LAGRANGIAN AND THE FERMIONIC COUPLINGS

We shall follow Refs. [13, 17], where a specific four-zero texture has been implemented for the Yukawa matrices within the 2HDM-III. This allows one to express the couplings of the neutral and charged Higgs bosons in terms of the fermion masses, Cabibbo-Kobayashi-Maskawa (CKM) mixing angles and certain dimensionless parameters, which are to be bounded by current experimental constraints. Thus, in order to derive the interactions of the charged Higgs boson, the Yukawa Lagrangian is written as follows:

$$\mathcal{L}_Y = Y_1^u \bar{Q}_L \tilde{\Phi}_1 u_R + Y_2^u \bar{Q}_L \tilde{\Phi}_2 u_R + Y_1^d \bar{Q}_L \Phi_1 d_R + Y_2^d \bar{Q}_L \Phi_2 d_R, \quad (1)$$

where $\Phi_{1,2} = (\phi_{1,2}^+, \phi_{1,2}^0)^T$ refer to the two Higgs doublets, $\tilde{\Phi}_{1,2} = i\sigma_2 \Phi_{1,2}^*$, Q_L denotes the left-handed fermion doublet, u_R and d_R are the right-handed fermions singlets and, finally, $Y_{1,2}^{u,d}$ denote the (3×3) Yukawa matrices. Similarly, one can write the corresponding Lagrangian for leptons.

After spontaneous EWSB and including the diagonalizing matrices for quarks and Higgs bosons², the interactions of the charged Higgs boson H^+ with quark pairs have the following form:

$$\begin{aligned} \mathcal{L}^{\bar{q}_i q_j H^+} = & \frac{g}{2\sqrt{2}M_W} \sum_{l=1}^3 \bar{u}_i \left\{ (V_{\text{CKM}})_{il} \left[\tan \beta m_{d_l} \delta_{lj} - \sec \beta \left(\frac{\sqrt{2}M_W}{g} \right) (\tilde{Y}_2^d)_{lj} \right] \right. \\ & + \left[\cot \beta m_{u_i} \delta_{il} - \csc \beta \left(\frac{\sqrt{2}M_W}{g} \right) (\tilde{Y}_1^u)_{il}^\dagger \right] (V_{\text{CKM}})_{lj} \\ & + (V_{\text{CKM}})_{il} \left[\tan \beta m_{d_l} \delta_{lj} - \sec \beta \left(\frac{\sqrt{2}M_W}{g} \right) (\tilde{Y}_2^d)_{lj} \right] \gamma^5 \\ & \left. - \left[\cot \beta m_{u_i} \delta_{il} - \csc \beta \left(\frac{\sqrt{2}M_W}{g} \right) (\tilde{Y}_1^u)_{il}^\dagger \right] (V_{\text{CKM}})_{lj} \gamma^5 \right\} d_j H^+, \end{aligned} \quad (2)$$

where V_{CKM} denotes the mixing matrices of the quark sector (and similarly for the leptons). The term proportional to δ_{ij} corresponds to the contribution that would arise within the 2HDM-II, while the terms proportional to $\tilde{Y}_2^d$ and $\tilde{Y}_1^u$ denote the new contributions from the 2HDM-III. These contributions, depend on the rotated matrices: $\tilde{Y}_n^q = O_q^T P_q Y_n^q P_q^\dagger O_q$ ($n = 1$ when $q = u$, and $n = 2$ when $q = d$), where O_q is the diagonalizing matrix, while P_q includes the phases of the Yukawa matrix. In order to evaluate $\tilde{Y}_n^q$ we shall consider that all Yukawa matrices have the

² The details of both diagonalizations are presented in Ref. [13].

four-Hermitic-texture form [15], and the quark masses have the same form, which are given by:

$$M^q = \begin{pmatrix} 0 & C_q & 0 \\ C_q^* & \tilde{B}_q & B_q \\ 0 & B_q^* & A_q \end{pmatrix} \quad (q = u, d). \quad (3)$$

To diagonalize these matrices, we use the matrices O_q and P_q , in the following way [15]:

$$\bar{M}^q = O_q^T P_q M^q P_q^\dagger O_q. \quad (4)$$

Then, one can derive a better approximation for the product $O_q^T P_q Y_n^q P_q^\dagger O_q$, expressing the rotated matrix $\tilde{Y}_n^q$, in the form

$$[\tilde{Y}_n^q]_{ij} = \frac{\sqrt{m_i^q m_j^q}}{v} [\tilde{\chi}_n^q]_{ij} = \frac{\sqrt{m_i^q m_j^q}}{v} [\chi_n^q]_{ij} e^{i\theta_{ij}^q}. \quad (5)$$

In order to perform our phenomenological study, we find it convenient to rewrite the Lagrangian given in Eq. (2) in terms of the coefficients $[\tilde{\chi}_n^q]_{ij}$, as follows:

$$\begin{aligned} \mathcal{L}^q = & \frac{g}{2\sqrt{2}M_W} \sum_{l=1}^3 \bar{u}_i \left\{ (V_{\text{CKM}})_{il} \left[\tan \beta m_{d_l} \delta_{lj} - \frac{\sec \beta}{\sqrt{2}} \sqrt{m_{d_l} m_{d_j}} \tilde{\chi}_{lj}^d \right] \right. \\ & + \left[\cot \beta m_{u_i} \delta_{il} - \frac{\csc \beta}{\sqrt{2}} \sqrt{m_{u_i} m_{u_l}} \tilde{\chi}_{il}^u \right] (V_{\text{CKM}})_{lj} \\ & + (V_{\text{CKM}})_{il} \left[\tan \beta m_{d_l} \delta_{lj} - \frac{\sec \beta}{\sqrt{2}} \sqrt{m_{d_l} m_{d_j}} \tilde{\chi}_{lj}^d \right] \gamma^5 \\ & \left. - \left[\cot \beta m_{u_i} \delta_{il} - \frac{\csc \beta}{\sqrt{2}} \sqrt{m_{u_i} m_{u_l}} \tilde{\chi}_{il}^u \right] (V_{\text{CKM}})_{lj} \gamma^5 \right\} d_j H^+, \end{aligned} \quad (6)$$

where we have redefined $[\tilde{\chi}_1^u]_{ij} = \tilde{\chi}_{ij}^u$ and $[\tilde{\chi}_2^d]_{ij} = \tilde{\chi}_{ij}^d$. Then, from Eq. (6), the couplings $\bar{u}_i d_j H^+$ and $u_i \bar{d}_j H^-$ are given by:

$$g_{H^+ \bar{u}_i d_j} = -\frac{ig}{2\sqrt{2}M_W} (S_{ij} + P_{ij} \gamma_5), \quad g_{H^- u_i \bar{d}_j} = -\frac{ig}{2\sqrt{2}M_W} (S_{ij} - P_{ij} \gamma_5), \quad (7)$$

where S_{ij} and P_{ij} are defined as:

$$\begin{aligned} S_{ij} = & \sum_{l=1}^3 \left\{ (V_{\text{CKM}})_{il} \left[\tan \beta m_{d_l} \delta_{lj} - \frac{\sec \beta}{\sqrt{2}} \sqrt{m_{d_l} m_{d_j}} \tilde{\chi}_{lj}^d \right] \right. \\ & \left. + \left[\cot \beta m_{u_i} \delta_{il} - \frac{\csc \beta}{\sqrt{2}} \sqrt{m_{u_i} m_{u_l}} \tilde{\chi}_{il}^u \right] (V_{\text{CKM}})_{lj} \right\}, \\ P_{ij} = & \sum_{l=1}^3 \left\{ (V_{\text{CKM}})_{il} \left[\tan \beta m_{d_l} \delta_{lj} - \frac{\sec \beta}{\sqrt{2}} \sqrt{m_{d_l} m_{d_j}} \tilde{\chi}_{lj}^d \right] \right. \\ & \left. - \left[\cot \beta m_{u_i} \delta_{il} - \frac{\csc \beta}{\sqrt{2}} \sqrt{m_{u_i} m_{u_l}} \tilde{\chi}_{il}^u \right] (V_{\text{CKM}})_{lj} \right\}. \end{aligned} \quad (8)$$

As it was discussed in Ref. [13], most low-energy processes imply weak bounds on the coefficients $\tilde{\chi}_{ij}^q$, which turn out to be of $O(1)$. However, some important constraints on $\tan\beta$ have started to appear, based on B -physics [31]. In order to discuss these results we find convenient to generalize the notation of Ref. [32] and define the couplings $\bar{u}_i d_j H^+$ and $u_i \bar{d}_j H^-$ in terms of the matrices X_{ij} , Y_{ij} and Z_{ij} (for leptons). In our case these matrices are given by:

$$\begin{aligned} X_{lj} &= \left[\tan\beta \delta_{lj} - \frac{\sec\beta}{\sqrt{2}} \sqrt{\frac{m_{d_j}}{m_{d_l}}} \tilde{\chi}_{lj}^d \right], \\ Y_{il} &= \left[\cot\beta \delta_{il} - \frac{\csc\beta}{\sqrt{2}} \sqrt{\frac{m_{u_l}}{m_{u_i}}} \tilde{\chi}_{il}^u \right]. \end{aligned} \quad (9)$$

where X_{lj} and Y_{il} are related with S_{ij} and P_{ij} defined in the Eq. (8) as follows:

$$\begin{aligned} S_{ij} &= \sum_{l=1}^3 (V_{\text{CKM}})_{il} m_{d_l} X_{lj} + m_{u_i} Y_{il} (V_{\text{CKM}})_{lj}, \\ P_{ij} &= \sum_{l=1}^3 (V_{\text{CKM}})_{il} m_{d_l} X_{lj} - m_{u_i} Y_{il} (V_{\text{CKM}})_{lj}. \end{aligned} \quad (10)$$

The 33 elements of these matrices reduce to the expressions for the parameters X, Y, Z ($= X_{33}, Y_{33}, Z_{33}$) used in Ref.[32]. Based on the analysis of $B \rightarrow X_s \gamma$ [32, 33], it is claimed that $X \leq 20$ and $Y \leq 1.7$ for $m_{H^+} > 250$ GeV, while for a lighter charged Higgs boson mass, $m_{H^+} \sim 180$ GeV, one gets $(X, Y) \leq (18, 0.5)$. In recently work we get the values of (X, Y) as a function of $\tan\beta$ within our model. Thus, we find the bounds: $|\chi_{33}^{u,d}| \lesssim 1$ for $0.1 < \tan\beta \leq 70$ [14]. Although in our model there are additional contributions (for instance from c -quarks, which are proportional to X_{23}), they are not relevant because the Wilson coefficients in the analysis of $B \rightarrow X_s \gamma$ are functions of m_c^2/M_W^2 or $m_c^2/m_{H^+}^2$ [34], that is, negligible when compared to the leading X_{33} effects, whose Wilson coefficients depend on m_t^2/M_W^2 or $m_t^2/m_{H^+}^2$. Other constraints on the charged Higgs mass and $\tan\beta$, can be obtained the anomalous magnetic momentum of the muon based on Δa_μ , the ρ parameter, as well as B -decays into the tau lepton, can be obtained [35, 36]. For instance, as can be read from Ref.[37], one has that the decay $B \rightarrow \tau \nu$, implies a constraint such that for $m_{H^+} = 200$ (300) GeV, values of $\tan\beta$ less than about 30 (50) are still allowed, within MSSM or THDM-II: However, these constraints can only be taken as estimates, as it is likely that they would be modified for THDM-III. In summary, we find that low energy constraints still allow to have $\tilde{\chi}_{ij}^q = O(1)^3$. On the other hand, the condition $\frac{\Gamma_{H^+}}{m_{H^+}} < \frac{1}{2}$ in the frame of the 2HDM-II implies $\frac{\Gamma_{H^+}}{m_{H^+}} \approx \frac{3G_F m_t^2}{4\sqrt{2}\pi \tan\beta^2}$ which leads to $0.3 \lesssim \tan\beta \lesssim 130$. However, in the 2HDM-III we have that

³ A more detailed analysis that includes the most recent data is underway [38, 39].

$\frac{\Gamma_{H^+}}{m_{H^+}} \approx \frac{3G_F m_t^2}{4\sqrt{2}\pi \tan \beta^2} \left(\frac{1}{1 - \frac{\tilde{\chi}_{33}^u}{\sqrt{2} \cos \beta}} \right)^2$, we have checked numerically that this leads to $0.08 < \tan \beta < 200$ when $|\tilde{\chi}_{33}^u| \approx 1$ and $0.3 < \tan \beta < 130$ as long as $|\tilde{\chi}_{33}^u| \rightarrow 0$ recovering the result for the case of the 2HDM-II [7, 40]. In this sense, if we consider the constraints imposed by the perturbativity bound, a portion of the low $\tan \beta$ appearing in some graphs would be excluded. However, we have decided to keep that range both to show the behaviour of the quantities of interest, and also to keep in mind that such criteria (perturbativity) should be taken as an order of magnitude constraint.

III. DECAY $H^+ \rightarrow W^+ \gamma$

In the calculation of the width for decay $H^+ \rightarrow W^+ \gamma$, we employ a nonlinear R_ξ -gauge, which leads to considerable simplifications due to the fact that some unphysical vertices are removed from the interaction Lagrangian [19]. We have shown that such a gauge not only reduces considerably the number of Feynman diagrams but also renders manifestly gauge-invariant and ultraviolet-finite amplitudes. Apart from emphasizing the advantages of using the nonlinear R_ξ -gauge, we will analyze the $H^+ \rightarrow W^+ \gamma$ decay in some scenarios which are still consistent with the most recent bounds discussed in the previous section.

A. The gauge fixing procedure

We now would like to comment on the gauge-fixing procedure which was used to simplify our calculation. To this end we introduce the following gauge-fixing functions [41]:

$$f^+ = \left(D_\mu^e + \frac{igs_W^2}{c_W} Z_\mu \right) W^{+\mu} - i\xi m_W G_W^+, \quad (11)$$

$$f^Z = \partial_\mu Z^\mu - \xi m_Z G_Z, \quad (12)$$

$$f^A = \partial_\mu A^\mu, \quad (13)$$

with D_μ^e the electromagnetic covariant derivative and ξ the gauge parameter. Note that f^+ is nonlinear and transforms covariantly under the electromagnetic gauge group. This gauge-fixing procedure is suited to remove the unphysical vertices $WG_W \gamma$ and $WG_W Z$, which arise in the Higgs kinetic-energy sector, and also modifies the Yang-Mills sector. One important result is that the expression for the $WW\gamma$ vertex satisfies a QED-like Ward identity, which turns out to be very useful in loop calculations. In particular, in the Feynman-t'Hooft gauge the Feynman rule for the $W_\rho^+(q)W_\nu^-(p)A_\mu(k)$ coupling can be written as

$$\Gamma_{\rho\nu\mu}^{WW\gamma} = -ie (g_{\mu\nu}(p-k+q)_\rho + g_{\nu\rho}(q-p)_\mu + g_{\mu\rho}(k-p-q)_\nu), \quad (14)$$

where all the momenta are incoming. It is easy to see that this expression satisfies a QED-like Ward identity. The remaining Feynman rules necessary for our calculation do not depend on the gauge fixing procedure. As far as the Yukawa couplings are concerned, we will concentrate on the type-III 2HDM.

B. Decays of the charged Higgs boson at tree level

The expressions for the charged Higgs boson decay widths $H^+ \rightarrow u_i \bar{d}_j$ are of the form:

$$\Gamma(H^+ \rightarrow u_i \bar{d}_j) = \frac{3g^2}{32\pi M_W^2 m_{H^+}^3} \lambda^{1/2}(m_{H^+}^2, m_{u_i}^2, m_{d_j}^2) \times \left(\frac{1}{2} \left[m_{H^+}^2 - m_{u_i}^2 - m_{d_j}^2 \right] (S_{ij}^2 + P_{ij}^2) - m_{u_i} m_{d_j} (S_{ij}^2 - P_{ij}^2) \right), \quad (15)$$

where λ is the usual kinematic factor $\lambda(a, b, c) = (a - b - c)^2 - 4bc$. When we replace $\tilde{\chi}_{ud} \rightarrow 0$, the formulae of the decays width become those of the 2HDM-II: see, e.g., Ref. [2]. Furthermore, the expressions for the charged Higgs boson decay widths of the bosonic modes remain the same as in the 2HDM-II. Recently [14], we have studied that the effect of the modified Higgs couplings typical of the 2HDM-III shows up clearly in the pattern of charged Higgs boson decays, which can be very different from the 2HDM-II case and thus enrich the possibilities to search for $H^\pm$ states at current (Tevatron) and future (LHC, ILC/CLIC) machines.

C. Decay $H^+ \rightarrow W^+ \gamma$ induced at one-loop level

In the nonlinear R_ξ -gauge, the decay $H^+ \rightarrow W^+ \gamma$ receives contributions from the Feynman diagrams shown in Fig. 1. As far as the fermionic sector is concerned, the main contribution comes from the third-generation quarks (t, b) , which induce three diagrams. However, we also considered the quarks contribution (c, b) and (t, s) because their coupling with the charged Higgs can be important [14], which induce six more diagram. Whereas in the bosonic sector there are contributions from the $(H^\pm, \phi^0)$ and $(W^\pm, \phi^0)$ pairs, with $\phi^0 = h^0$ or H^0 . We would like to emphasize that the nonlinear R_ξ -gauge considerably simplifies the calculation of the decay $H^+ \rightarrow W^+ \gamma$. First of all, the removal of the unphysical vertex $W^\pm G_W^\mp \gamma$, and the tadpole graphs vanish [19]. Apart from these simplifications, we will be able to group the Feynman diagrams into subsets which separately yield a manifestly gauge-invariant amplitude free of ultraviolet singularities. Once the amplitude for every Feynman diagram shown in Fig. 1 was written down, the Passarino-Veltman method [42] was applied to express it in terms of scalar integrals, which are suitable for




FIG. 1: Feynman diagrams contributing to the $H^+ \rightarrow W^+ \gamma$ decay in the nonlinear R_ξ -gauge. ϕ^0 stands for h^0 and H^0 , and X^+ for H^+ and G_W^+ .

numerical evaluation [43]. The full amplitude for the decay $H^+(p) \rightarrow W^+(q)\gamma(k)$ can be written as

$$\mathcal{M}(H^+ \rightarrow W^+ \gamma) = \frac{\alpha^{3/2}}{2\sqrt{\pi}m_W s_w^2} \left(G (q \cdot k g_{\mu\nu} - k_\nu q_\mu) - i H \epsilon_{\mu\nu\alpha\beta} k^\alpha q^\beta \right) \epsilon^\mu(k) \epsilon^\nu(q), \quad (16)$$

which is manifestly gauge invariant. The H function only receives contributions from the quarks:

$$H = \frac{i}{\Delta_{(H^\pm, W)}} \sum_{i,j}^3 \left(3 \lambda_{ij} \Delta B_{(H^\pm, W, d_j, u_i)} + \Delta_{(H^\pm, W)} \left(\lambda_{ij}^d C_{(d_j, u_i)} - 2 \lambda_{ij}^u C_{(u_i, d_j)} \right) \right), \quad (17)$$

where

$$\lambda_{ij} = -\frac{1}{2} \left(m_{u_i} (P_{ij} + S_{ij}) - m_{d_j} (P_{ij} - S_{ij}) \right), \quad (18)$$

$$\lambda_{ij}^u = \frac{1}{2} m_{u_i} (P_{ij} + S_{ij}), \quad (19)$$

$$\lambda_{ij}^d = \frac{1}{2} m_{d_i} (P_{ij} - S_{ij}), \quad (20)$$

here, we have introduced the shorthand notation $\Delta_{(a,b)} \equiv m_a^2 - m_b^2$, $\Delta B_{(a,b,c,d)} = B_0(m_a^2, m_c^2, m_d^2) - B_0(m_b^2, m_c^2, m_d^2)$,⁴ and $C_{(a,b)} \equiv C_0(m_{H^\pm}^2, m_W^2, 0, m_a^2, m_b^2, m_a^2)$, where B_0 and C_0 stand for Passarino-Veltman scalar integrals. From the $\Delta B_{(a,b,c,d)}$ definition, it is clear that the H function is ultraviolet finite. Again, when we replace $\tilde{\chi}_{ud} \rightarrow 0$, the formulae of the decays width become those of the 2HDM-II [19]. As for G , it can be written as

$$G = \frac{1}{2 m_{H^\pm}^2 \Delta_{(H^\pm, W)}^2} \left(G_{u_i d_j} + \sum_{\phi^0=H^0, h^0} \left(G_{H^\pm \phi^0} + G_{G_W^\pm \phi^0} + G_{W^\pm \phi^0} \right) \right), \quad (21)$$

⁴ When $a = 0$, m_a must be substituted by 0.

where G_{AB} stands for the contribution of the (A, B) pair:

$$\begin{aligned}
G_{u_i d_j} = & \sum_{ij}^3 m_{H^\pm}^2 \left(\left(3 \Delta_{(u_i, d_j)} + m_W^2 \right) \lambda_{ij} + 3 \sigma_{ij} \Delta_{(H^\pm, W)} \right) \Delta B_{(H^\pm, W, d_j, u_i)} \\
& - 3 \Delta_{(u_i, d_j)} \Delta_{(H^\pm, W)} \lambda_{ij} \Delta B_{(0, H^\pm, d_j, u_i)} + m_{H^\pm}^2 \Delta_{(H^\pm, W)} \left(\lambda_{ij} - (2 m_{d_j}^2 \lambda_{ij} - \lambda_{ij}^d \Delta_{(H^\pm, W)}) C_{(d_j, u_i)} \right. \\
& \left. + (2 m_{u_i}^2 \lambda_{ij} - \lambda_{ij}^u \Delta_{(H^\pm, W)}) C_{(u_i, d_j)} \right), \tag{22}
\end{aligned}$$

where

$$\sigma_{ij} = \frac{1}{2} \left(m_{u_i} (P_{ij} + S_{ij}) + m_{d_j} (P_{ij} - S_{ij}) \right), \tag{23}$$

When one takes $\chi'_{ij}s \rightarrow 0$, the formulas for the function $G_{u_i d_j}$ reduce to the 2HDM-II case, see, e. g. [19].

$$\begin{aligned}
G_{H^\pm h^0} = & \lambda_{h^0 H^\pm} \left(\Delta_{(H^\pm, h^0)} \Delta_{(H^\pm, W)} \delta B_{(0, W, H^\pm, h^0)} - (2 m_{H^\pm}^2 \Delta_{(H^\pm, h^0)} + m_{h^0}^2 m_W^2) \delta B_{(H^\pm, W, H^\pm, h^0)} \right. \\
& \left. - m_{H^\pm}^2 \Delta_{(H^\pm, W)} (1 + 2 m_{H^\pm}^2 C_{(H^\pm, h^0)}) \right), \tag{24}
\end{aligned}$$

$$\begin{aligned}
G_{G_W^\pm h^0} = & \lambda_{G_W^\pm h^0} \left((m_W^2 \Delta_{(h^0, W)} + m_{H^\pm}^2 (3 m_W^2 - 2 m_{h^0}^2)) \delta B_{(H^\pm, W, h^0, W)} + \Delta_{(H^\pm, h^0)} \Delta_{(h^0, W)} \delta B_{(0, W, h^0, W)} \right. \\
& \left. + m_{H^\pm}^2 \Delta_{(H^\pm, W)} (1 + 2 m_W^2 C_{(W, h^0)}) \right), \tag{25}
\end{aligned}$$

and

$$\begin{aligned}
G_{W^\pm H^0} = & \lambda_{W^\pm H^0} \left((m_W^2 \Delta_{(h^0, W)} + m_{H^\pm}^2 (7 m_W^2 - 2 m_{h^0}^2 - 4 m_{H^\pm}^2)) \delta B_{(H^\pm, W, H^\pm, h^0)} \right. \\
& + \Delta_{(H^\pm, W)} \Delta_{(h^0, W)} \delta B_{(0, W, H^\pm, h^0)} \\
& \left. + m_{H^\pm}^2 \Delta_{(H^\pm, W)} \left(1 - 2 (2 \Delta_{(H^\pm, W)} - m_W^2) C_{(m_W^2, m_{h^0}^2)} \right) \right), \tag{26}
\end{aligned}$$

with

$$\lambda_{H^\pm h^0} = c_{\alpha-\beta} \left(2 s_{\alpha-\beta} m_{H^\pm}^2 + \frac{c_{\alpha+\beta}}{c_\beta^2 s_\beta^2} \mu_{12}^2 + \left(s_\alpha s_\beta t_\beta - \frac{c_\beta c_\alpha}{t_\beta} \right) m_{h^0}^2 \right), \tag{27}$$

$$\lambda_{G_W^\pm h^0} = \frac{s_{2(\alpha-\beta)} \Delta_{(H^\pm, h^0)}}{2}, \tag{28}$$

and

$$\lambda_{W^\pm H^0} = -\frac{m_W^2 s_{2(\alpha-\beta)}}{2}. \tag{29}$$

Finally, the contribution of the heaviest CP-even scalar boson H^0 is obtained of the lightest one, once the substitutions $m_{h^0} \rightarrow m_{H^0}$ and $\alpha \rightarrow \alpha - \pi/2$ are done.

It is also evident that the partial amplitudes induced by the pairs (t, b) , (t, s) , (c, b) , $(\phi^0, W^\pm)$, $(\phi^0, H^\pm)$, and $(\phi^0, G_W^\pm)$ are gauge invariant and ultraviolet finite on their own. Again, this is to be contrasted with the situation arising in the linear R_ξ -gauge, where showing gauge invariance is somewhat cumbersome, and the cancellation of ultraviolet divergences in the bosonic sector is achieved only after adding up all the Feynman diagrams [19].

Once Eq. (16) is squared and the spins of the final particles are summed over, the decay width can be written as

$$\Gamma(H^+ \rightarrow W^+ \gamma) = \frac{\alpha^3 \Delta_{(H^\pm, W)}^3}{2^7 \pi^2 s_W^4 m_W^2 m_{H^\pm}^3} (|G|^2 + |H|^2). \quad (30)$$

We will evaluate this decay width for some values of the parameters of the model.

D. Numerical results and discussion

Let us now discuss the decay modes of the charged Higgs boson within our model. Hereafter, we shall refer to two benchmark scenarios, namely. (i) **Scenario A**: $\tilde{\chi}_{ij}^u = -1$, $\tilde{\chi}_{ij}^d = -1$; (ii) **Scenario B**: $\tilde{\chi}_{ij}^u = 1$, $\tilde{\chi}_{ij}^d = 1$. We have performed the numerical analysis of charged Higgs boson decays by taking $\tan \beta = 0.3, 0.5, 1, 10$ and varying the charged Higgs boson mass within the interval $100 \text{ GeV} \leq m_{H^\pm} \leq 800 \text{ GeV}$, further fixing $m_{h^0} = 120 \text{ GeV}$, $m_{A^0} = 300 \text{ GeV}$ and the mixing angle at $\alpha = \pi/2$. Then the results for the Branching Ratios (BRs) are shown in Fig. 2, and have the following characteristics.

Scenario A. In Fig. 2(a) we present the BRs for the channels $H^+ \rightarrow W^+ \gamma$, $H^+ \rightarrow t\bar{b}$, $c\bar{b}$, $t\bar{s}$, $\tau^+ \nu_\tau$, $W^+ h^0$, $W^+ A^0$ as a function of m_{H^+} , for $\tan \beta = 0.3$ and fixing $m_{h^0} = 120 \text{ GeV}$, $m_{A^0} = 300 \text{ GeV}$ and the mixing angle $\alpha = \pi/2$. When $m_{H^+} < 175 \text{ GeV}$, we can see that the dominant decay of the charged Higgs boson is via the mode $\tau\bar{\nu}$, with $\text{BR}(H_i^+ \rightarrow \tau\bar{\nu}) \approx 1$, and the same importance we have the mode $c\bar{b}$. For the case $170 \text{ GeV} < m_{H^+} < 180 \text{ GeV}$ the mode $t\bar{s}$ is relevant. In this range, the mode $W^+ \gamma$ is competitive and becomes of order 10^{-1} , which it is also very different from the 2HDM-II and becomes an interesting phenomenological consequence of the 2HDM-III. We can also observe that, for $m_{H^+} > 180 \text{ GeV}$, the decay mode $t\bar{b}$ is dominant (as in the 2HDM-II). In this range mass of the charged Higgs, the channel $H^+ \rightarrow W^+ \gamma$ has a BR $\sim 10^{-4}$ to $\sim 10^{-6}$. Now, from Fig. 2(b), where $\tan \beta = 0.5$, we find that the dominant decay mode is into $\tau^+ \nu_\tau$ for the range $m_{H^+} < 175 \text{ GeV}$, again for $175 \text{ GeV} < m_{H^+} < 180 \text{ GeV}$ the mode $t\bar{s}$ is the leading one, but for $180 \text{ GeV} < m_{H^+}$, the decay channel $t\bar{b}$ becomes relevant, whereas for the range $170 \text{ GeV} < m_{H^+} < 180 \text{ GeV}$ the mode $W^+ \gamma$ has a BR of order 10^{-2} . Again, when $m_{H^+} > 180 \text{ GeV}$ the




FIG. 2: The figure shows the BRs of the H^+ decaying into the principal modes in Scenario A, taking $\tilde{\chi}_{ij}^u = -1$, $\tilde{\chi}_{ij}^d = -1$, $m_{h^0} = 120$ GeV, $m_{A^0} = 120$ GeV and $\alpha = \pi/2$ for: (a) $\tan \beta = 0.3$, (b) $\tan \beta = 0.5$, (c) $\tan \beta = 1$, (d) $\tan \beta = 10$. The lines in each graph correspond to: ($W\gamma$ line) $\text{BR}(H^+ \rightarrow W^+\gamma)$, (tb line) $\text{BR}(H^+ \rightarrow t\bar{b})$, (cb line) $\text{BR}(H^+ \rightarrow c\bar{b})$, (ts line) $\text{BR}(H^+ \rightarrow t\bar{s})$, ($\tau\nu$ line) $\text{BR}(H^+ \rightarrow \tau^+\nu_\tau)$, (Wh line) $\text{BR}(H^+ \rightarrow W^+h^0)$, (WA line) $\text{BR}(H^+ \rightarrow W^+A^0)$.

$\text{BR}(H^+ \rightarrow W^+\gamma)$ is of order 10^{-4} to 10^{-7} . Then, see Fig. 2(c), for the case with $\tan \beta = 1$ one gets that $\text{BR}(H^+ \rightarrow \tau^+\nu_\tau) \approx 1$ when $m_{H^+} < 175$ GeV, while in the range $175 \text{ GeV} < m_{H^+} < 180$ GeV the mode $t\bar{s}$ is relevant and the $\text{BR}(H^+ \rightarrow W^+\gamma)$ even could be important with an order 10^{-3} . However for $180 \text{ GeV} < m_{H^+}$, the dominant decay of the charged Higgs boson is the mode $t\bar{b}$. For $\tan \beta = 10$, we show in plot Fig. 2(d) that the dominant decay of the charged Higgs boson is the mode $\tau^+\nu_\tau$, when $m_{H^+} < 180$ GeV, but that, for $180 \text{ GeV} < m_{H^+} < 250$ GeV, the decay channel $t\bar{b}$ becomes the leading one, whereas for the range $250 \text{ GeV} < m_{H^+}$, the mode W^+h^0 is relevant. In this case the BR of the decay channel $W^+\gamma$ is unimportant.




FIG. 3: The same as in Fig. 2 but taking $\tilde{\chi}_{ij}^u = 1$, $\tilde{\chi}_{ij}^d = 1$ (Scenario B).

Scenario B. In Fig. 3(a) we present the BRs for the channels $H^+ \rightarrow W^+\gamma, t\bar{b}, c\bar{b}, t\bar{s}, \tau^+\nu_\tau, W^+h^0, W^+A^0$ as a function of m_{H^+} , for $\tan\beta = 0.3$ and fixing $m_{h^0} = 120$ GeV, $m_{A^0} = 300$ GeV and the mixing angle $\alpha = \pi/2$. When $m_{H^+} < 175$ GeV, we can see that the dominant decay of the charged Higgs boson is via the mode $\tau^+\nu_\tau$, with $\text{BR}(H_i^+ \rightarrow \tau^+\nu_\tau) \approx 1$. For the case $175 \text{ GeV} < m_{H^+} < 180$ GeV the mode $t\bar{s}$ is dominant, whereas for the range $170 \text{ GeV} < m_{H^+} < 180$ GeV the mode $W^+\gamma$ has a BR of order 10^{-2} . We can also observe that, for $m_{H^+} > 180$ GeV, the decay mode $t\bar{b}$ is dominant (as in the 2HDM-II). Now, from Fig. 3(b), where $\tan\beta = 0.5$, we find that the dominant decay mode is into $\tau^+\nu_\tau$ for the range $m_{H^+} < 175$ GeV, again for $175 \text{ GeV} < m_{H^+} < 180$ GeV the mode $t\bar{s}$ is the leading one, while the decay $H^+ \rightarrow W^+\gamma$ induced at one-loop level has a BR relatively large of order 10^{-2} . However for $180 \text{ GeV} < m_{H^+}$, the dominant decay of the charged Higgs boson is the mode $t\bar{b}$. Then, see Fig. 3(c), for the case with $\tan\beta = 1$ one gets that $\text{BR}(H^+ \rightarrow \tau^+\nu_\tau) \approx 1$ when $m_{H^+} < 175$ GeV, while in the range $175 \text{ GeV} < m_{H^+} < 180$ GeV the

mode $t\bar{s}$ is relevant and the $BR(H^+ \rightarrow W^+\gamma)$ even is of order 10^{-4} . For $180 \text{ GeV} < m_{H^+} < 600 \text{ GeV}$, the decay channel W^+h^0 becomes relevant, whereas for the range $600 \text{ GeV} < m_{H^+}$ the mode W^+A^0 is dominant. It is convenient to mention that this sub-scenario is special for the mode $t\bar{b}$, because its decay width is zero at the tree-level, since the CKM contribution is canceled exactly with the terms of the four-zero texture implemented for the Yukawa coupling of the 2HDM-III. For $\tan\beta = 10$, we show in plot Fig. 3(d) that the dominant decay of the charged Higgs boson is the mode $\tau^+\nu_\tau$, when $m_{H^+} < 180 \text{ GeV}$, but for $180 \text{ GeV} < m_{H^+} < 250 \text{ GeV}$, the decay channel $t\bar{b}$ becomes the leading one, whereas for the range $250 \text{ GeV} < m_{H^+}$, the mode W^+h^0 is again dominant. The decay $H^+ \rightarrow W^+\gamma$ has a BR of order 10^{-5} to 10^{-7} , which could be irrelevant channel decay for LHC.

In order to cover further the Higgs sector in our analysis, it is appropriate to also mention how the previous results change with m_{h^0} , m_{A^0} and α . In general the behavior of the decay modes of the charged Higgs boson is similar to the cases presented above, except for the decay channel Wh^0 , which can be important or unimportant. For $\alpha = 0$, this mode has $BR < 10^{-3}$ when $\tan\beta$ is large. However, for $\tan\beta < 1$, it becomes the dominant one. In the case $\alpha = \beta$, the decay channel Wh^0 can be the dominant mode, with a BR of $O(1)$.

As a general lesson from this section, we notice a distinctive features of our 2HDM-III, namely that the decay modes $W^+\gamma$ becomes very large, with a $BR = 2 \times 10^{-1}$ or 10^{-2} for some of the scenarios considered, with $\tan\beta < 2$. We also want to compare the general behavior of these decay modes, within 2HDM-III with the 2HDM-II case. In order to compare the 2HDM-III results with those in the 2HDM-II, we show in Fig. 4 the $BR(H^+ \rightarrow W^+\gamma)_{\text{III}}$ in the scenario A and B from 2HDM-III, as well as $BR(H^+ \rightarrow W^+\gamma)_{\text{II}}$ from 2HDM-II *vs.* m_{H^+} , taking again $\tan\beta = 0.3, 0.5, 1, 10$. We observe that the mode $W^+\gamma$ is important when $170 \text{ GeV} < m_{H^+} < 180 \text{ GeV}$ and for $0.1 \leq \tan\beta \leq 1$, taking $\tilde{\chi}_{ij}^{u,d} = -1$. Thus, we find that the effect of the modified Higgs couplings typical of the 2HDM-III shows up clearly in the pattern of charged Higgs boson decay $H^+ \rightarrow W^+\gamma$, which can be very different from the 2HDM-II case, and thus enrich the possibilities to search for $H^\pm$ states at current (Tevatron) and future (LHC, ILC/CLIC) machines.

IV. EVENTS RATES AT LHC FROM THE DECAY $H^+ \rightarrow W^+\gamma$

The production of charged Higgs bosons at hadron colliders has been evaluated in [23, 24] (also for the Superconducting Super Collider, SSC) and more recently [14, 25] (for the LHC) literature,




FIG. 4: The figure shows the $BR(H^+ \rightarrow W^+ \gamma)_{\text{III}}$, $BR(H^+ \rightarrow W^+ \gamma)_{\text{II}}$ vs. m_{H^+} , taking $\tan \beta = 0.3, 0.5, 1, 10$ for $\tilde{\chi}_{ij}^{u,d} = -1$ (dots), $\tilde{\chi}_{ij}^{u,d} = 0$ or case 2HDM-II (solid line), $\tilde{\chi}_{ij}^{u,d} = 1$ (dashes).

mainly for the 2HDM-II and its SUSY realization (i.e., the MSSM). In these two scenarios, when kinematically allowed, the top quark decay channel $t \rightarrow bH^+$ is the dominant $H^\pm$ production mechanism. Instead, above the threshold for such a decay, the dominant $H^\pm$ production reaction is gluon-gluon fusion into a 3-body final state, i.e., $gg \rightarrow tbH^\pm$ ⁵. Both processes depend on the coupling $H^- t \bar{b}$ and are therefore sensitive to the modifications that arise in the 2HDM-III for this vertex. However, detection of the final state will depend on the charged Higgs boson decay mode, which could include a complicated final state, that could in turn be difficult to reconstruct. For these reasons, it is very important to look for other production channels, which may be easier to reconstruct. In this regard, the s -channel production of charged Higgs bosons, through the mechanism of $c\bar{b}$ -fusion, could help to make more viable the detection of several charged Higgs

⁵ In fact, these two mechanisms are intimately related, see below.

boson decay channels [30].

A. Evaluation of event rates from direct production of charged Higgs bosons at the LHC

The $H^\pm \bar{q}q'$ vertex with large flavor mixing coupling, that arises in the 2HDM-III, enables the possibility of studying the production of charged Higgs boson via the s -channel production mechanism, $c\bar{b} \rightarrow H^+ + \text{c.c.}$ This process was discussed first in Ref. [30] and recently by [14] in the context of the 2HDM-III.

To illustrate the type of charged Higgs signatures that have the potential to be detectable at the LHC in the 2HDM-III, we show in the Table I the event rates of charged Higgs boson through the channel $c\bar{b} \rightarrow H^+ + \text{c.c.}$, alongside the corresponding production cross sections (σ 's) and relevant $\text{BR}(H^+ \rightarrow W^+\gamma)$, for a combination of masses, $\tan\beta$ and specific 2HDM-III parameters amongst those used in the previous sections (assuming $m_{h^0} = 120$ GeV, $m_{A^0} = 300$ GeV and the mixing angle at $\alpha = \pi/2$ throughout). In particular, we focus on those cases where the charged Higgs boson mass is above the threshold for $t \rightarrow bH^+$, for two reasons. On the one hand, the scope of the LHC in accessing $t \rightarrow bH^+$ decays has been established in a rather model independent way. On the other hand, we have dealt at length with the corresponding BRs in section III. (As default, we also assume an integrated luminosity of 10^5 pb^{-1} .) In all cases we use the results of our previously work [14].

To illustrate these results, let us comment on one case within the scenario B, because is the case most conservative. From Table I, we can see that for Scenario B, with $(\tilde{\chi}_{ij}^u = 1, \tilde{\chi}_{ij}^d = 1)$ and $\tan\beta = 0.3$, we have that the $H^\pm$ is heavier than $m_t - m_b$, as we take a mass $m_{H^+} = 200$ GeV, thus precluding top decay contributions, so that in this case $\sigma(pp \rightarrow H^+X) \approx 2.2 \times 10^{-2}$ pb, and including the decay $H^+ \rightarrow W^+\gamma$, in the final state we set 42 events. Even for charged Higgs masses as long as 300 GeV, we can still obtain 4 events. The most promising rate for decay $H^+ \rightarrow W^+\gamma$, is for $m_{H^+} = 200$ and $\tan\beta = 1$, when we get 63 events.

B. Indirect production of charged Higgs bosons at the LHC

We have found that, in some of the 2HDM-III scenarios envisaged here, light charged Higgs bosons could exist that have not been excluded by current experimental bounds, chiefly from LEP2 and Tevatron. Their discovery potential should therefore be studied in view of the upcoming LHC and we shall then turn our attention now to presenting the corresponding hadro-production cross

TABLE I: Summary of LHC event rates for some parameter combinations within Scenarios B with an integrated luminosity of 10^5 pb^{-1} , for the signal $H^+ \rightarrow W^+ \gamma$, through the channel $c\bar{b} \rightarrow H^+ + \text{c.c.}$

$(\tilde{\chi}_{ij}^u, \tilde{\chi}_{ij}^d)$	$\tan \beta$	m_{H^+} in GeV	$\sigma(pp \rightarrow H^+ + X)$ in pb	$BR(H^+ \rightarrow W^+ \gamma)$	Nr. Events
(1,1)	0.3	200	2.1×10^2	2×10^{-6}	42
(1,1)	0.3	300	3.1×10	1.1×10^{-6}	4
(1,1)	1	200	3	2.1×10^{-4}	63
(1,1)	1	300	0.98	4.2×10^{-7}	0
(1,1)	10	200	1.1	9.8×10^{-5}	10
(1,1)	10	300	5.2×10^{-1}	1.1×10^{-6}	0

sections via an indirect channel, i.e., other than as secondary products in (anti)top quark decays and via $c\bar{b}$ -fusion, considered previously.

In previously work [14], we evaluate the $q\bar{q}, gg \rightarrow t\bar{b}H^- + \text{c.c.}$ cross sections. We use HERWIG version 6.510 in default configuration, by onsetting the subprocess `IPROC = 3839`, wherein we have overwritten the default MSSM/2HDM couplings and masses with those pertaining to the 2HDM-III: see Eqs. (7)–(8).

Now, let discuss again the scenario B. From Table II, we can see that for this scenario, with $(\tilde{\chi}_{ij}^u = 1, \tilde{\chi}_{ij}^d = 1)$ and $\tan \beta = 0.3$, we have that the $H^\pm$ is heavier than $m_t - m_b$, as we take a mass $m_{H^+} = 200 \text{ GeV}$, thus precluding top decay contributions, so that in this case $\sigma(pp \rightarrow t\bar{b}H^-) \approx 25.8 \text{ pb}$, while the signal $H^+ \rightarrow W^+ \gamma$ give a number of events of 5. When the mass of the charged Higgs take the value of 300 GeV, we can obtain only 1 event. One can see that for $m_{H^+} = 200$ and $\tan \beta = 1$, we get 48 events. However, the case most promising rate for decay $H^+ \rightarrow W^+ \gamma$ is when $m_{H^+} = 200$ and $\tan \beta = 10$, we get the spectacular 122 events, which could be important for the LHC collider experiments.

Altogether, by comparing the $q\bar{q}, gg \rightarrow t\bar{b}H^- + \text{c.c.}$ cross sections herein with, e.g., those of the MSSM in [6] or the 2HDM in [44, 45], it is clear that the 2HDM-III rates can be very large and thus the discovery potential in ATLAS and CMS can be substantial, particularly for a very light $H^\pm$, which may pertain to our 2HDM-III but not the MSSM or 2HDM-II. However, it is only by combining the production rates of this section with the decay ones of the previous ones that actual event numbers at the LHC can be predicted.

V. CONCLUSIONS

We have discussed the implications of assuming a four-zero Yukawa for the properties of the charged Higgs boson, within the context of a 2HDM-III. In particular, we have presented a detailed

TABLE II: Summary of LHC event rates for some parameter combinations within Scenarios B with for an integrated luminosity of 10^5 pb^{-1} , for $H^+ \rightarrow W^+\gamma$ signature, through the channel $q\bar{q}, gg \rightarrow \bar{t}bH^+ + \text{c.c.}$

$(\tilde{\chi}_{ij}^u, \tilde{\chi}_{ij}^d)$	$\tan \beta$	m_{H^+} in GeV	$\sigma(pp \rightarrow H^+ \bar{t}b)$ in pb	$\text{BR}(H^+ \rightarrow W^+\gamma)$	Nr. Events
(1,1)	0.3	200	25.8	2×10^{-6}	5
(1,1)	0.3	300	10.1	1.1×10^{-6}	1
(1,1)	1	200	2.3	2.1×10^{-4}	48
(1,1)	1	300	0.79	4.2×10^{-7}	0
(1,1)	10	200	12.5	9.8×10^{-5}	122
(1,1)	10	300	0.48	1.1×10^{-6}	0

discussion of the charged Higgs boson couplings to heavy fermions and the resulting implications for the decay $H^+ \rightarrow W^+\gamma$, induced at one-loop level. The latter clearly reflect the different coupling structure of the 2HDM-III, e.g., with respect to the 2HDM-II, so that one has at disposal more possibilities to search for $H^\pm$ states at current and future colliders, enabling us to distinguish between different models of EWSB. We have then concentrated our analysis to the case of the LHC and showed that the production rates of charged Higgs bosons at the LHC is sensitive to the modifications of the Higgs boson couplings. We have employed the results of the s -channel production of $H^\pm$ through $c\bar{b}$ -fusion and the multibody final state induced by gg -fusion and $q\bar{q}$ -annihilation. Finally, we have determined the number of events for the most promising LHC signatures for the decay $H^+ \rightarrow W^+\gamma$ within 2HDM-III, for both $c\bar{b} \rightarrow H^+ + \text{c.c.}$ and $q\bar{q} \rightarrow \bar{t}bH^+ + \text{c.c.}$ scatterings (the latter affording larger rates than the former). Armed with these results, we are now in a position to carry out a detailed study of signal and background rates, in order to determine the precise detectability level of each signature. However, this is beyond the scope of present work and will be the subject of a future publication.

Acknowledgements

We thank to Lorenzo Díaz-Cruz and Alfonso Rosado for useful discussions. This work was supported in part by CONACyT and SNI (México). J.H-S. thanks in particular CONACyT (México) for the grant J50027-F and SEP (México) by the grant PROMEP/103.5/08/1640. R.N-P. acknowledges the Institute of Physics BUAP for a warm hospitality and also financial support by

-
- [1] S. L. Glashow, Nucl. Phys. **22**, 579 (1961); S. Weinberg, Phys. Rev. Lett. **19**, 1264 (1967); A. Salam, Proc. 8th NOBEL Symposium, ed. N. Svartholm (Almqvist and Wiksell, Stockholm, 1968), p. 367.
 - [2] S. Dawson *et al.*, *The Higgs Hunter's Guide*, 2nd ed., Frontiers in Physics Vol. **80** (Addison-Wesley, Reading MA, 1990).
 - [3] M. Carena *et al.*, Report of the Tevatron Higgs working group; FERMILAB-CONF-00-279-T; hep-ph/0010338; C. Balazs *et al.*, Phys. Rev. **D59**, 055016 (1999).
 - [4] J.L. Díaz-Cruz *et al.*, Phys. Rev. Lett. **80**, 4641 (1998).
 - [5] J. Lorenzo Diaz Cruz, AIP Conf. Proc. **1026**, 30 (2008).
 - [6] A. Djouadi, Phys. Rept. **459**, 1 (2008).
 - [7] V. D. Barger, J. L. Hewett and R. J. N. Phillips, Phys. Rev. D **41**, 3421 (1990).
 - [8] See for instance: B. Dobrescu, Phys. Rev. **D63**, 015004 (2001). See also, recent work on Little Higgs models: N. Arkani-Hamed, A. G. Cohen, E. Katz and A. E. Nelson, JHEP **0207**, 034 (2002). And for AdS/CFT Higgs models: R. Contino, Y. Nomura and A. Pomarol, Nucl. Phys. B **671**, 148 (2003); A. Aranda, J. L. Diaz-Cruz, J. Hernandez-Sanchez and R. Noriega-Papaqui, Phys. Lett. B **658**, 57 (2007)
 - [9] J. Brau *et al.* [ILC Collaboration], arXiv:0712.1950 [physics.acc-ph]; A. Djouadi, J. Lykken, K. Monig, Y. Okada, M.J. Oreglia and S. Yamashita, arXiv:0709.1893 [hep-ph]; T. Behnke *et al.* [ILC Collaboration], arXiv:0712.2356 [physics.ins-det].
 - [10] G. Guignard (editor) [The CLIC Study Team], preprint CERN-2000-008 (2000).
 - [11] S. Kanemura, S. Moretti, Y. Mukai, R. Santos and K. Yagyu, arXiv:0901.0204 [hep-ph].
 - [12] K. S. Babu and C. F. Kolda, Phys. Lett. B **451**, 77 (1999).
 - [13] J. L. Diaz-Cruz, R. Noriega-Papaqui and A. Rosado, Phys. Rev. D **71**, 015014 (2005).
 - [14] J. L. Diaz-Cruz, J. Hernandez-Sanchez, S. Moretti, R. Noriega-Papaqui and A. Rosado, Phys. Rev. D **79**, 095025 (2009) [arXiv:0902.4490 [hep-ph]].
 - [15] H. Fritzsch and Z. Z. Xing, Phys. Lett. B **555** (2003) 63.
 - [16] T. P. Cheng and M. Sher, Phys. Rev. D **35**, 3484 (1987).
 - [17] J. L. Diaz-Cruz, R. Noriega-Papaqui and A. Rosado, Phys. Rev. D **69**, 095002 (2004).
 - [18] J. L. Díaz-Cruz and M.A. Pérez, Phys. Rev. D **33**, 273 (1986); J. Gunion, G. Kane and J. Wudka, Nucl. Phys. B **299**, 231 (1988); A. Mendez and A. Pomarol, Nucl. Phys. B **349**, 369 (1991); E. Barradas *et al.*, Phys. Rev. D **53**, 1678 (1996); M. Capdequi Peyranere, H. E. Haber and P. Irulegui, Phys. Rev. D **44**, 191 (1991); S. Moretti and W. J. Stirling, Phys. Lett. B **347**, 291 (1995) [Erratum-ibid. B **366**, 451 (1996)]; A. Djouadi, J. Kalinowski and P. M. Zerwas, Z. Phys. C **70**, 435 (1996); S. Kanemura, Phys. Rev. D **61**, 095001 (2000); E. Asakawa, S. Kanemura and J. Kanzaki, Phys. Rev. D **75**, 075022

- (2007); A. Arhrib, R. Benbrik and M. Chabab, J. Phys. G **34**, 907 (2007); A. Arhrib, R. Benbrik and M. Chabab, Phys. Lett. B **644**, 248 (2007).
- [19] J. Hernandez-Sanchez, M. A. Perez, G. Tavares-Velasco and J. J. Toscano, Phys. Rev. D **69**, 095008 (2004) [arXiv:hep-ph/0402284].
- [20] J. L. Díaz-Cruz, J. Hernández-Sánchez and J.J. Toscano, Phys. Lett. B **512**, 339 (2001).
- [21] J. L. Diaz-Cruz, J. Hernandez-Sanchez, S. Moretti and A. Rosado, Phys. Rev. D **77**, 035007 (2008) J. L. Diaz-Cruz, J. Hernandez-Sanchez, S. Moretti and A. Rosado, AIP Conf. Proc. **1026**, 138 (2008).
- [22] E. Barradas-Guevara, O. Félix-Beltrán, J. Hernández-Sánchez and A. Rosado, Phys. Rev. D **71**, 073004 (2005).
- [23] M. A. Pérez and A. Rosado, Phys. Rev. D **30**, 228 (1984); J. Gunion *et al.*, Nucl. Phys. B **294**, 621 (1987); J. F. Gunion, Phys. Lett. B **322**, 125 (1994).
- [24] J. L. Díaz-Cruz and O.A. Sampayo, Phys. Rev. D **50**, 6820 (1994).
- [25] S. Moretti and K. Odagiri, Phys. Rev. D **55**, 5627 (1997); Phys. Rev. D **59**, 055008 (1999); F. Borzumati, J.-L. Kneur and N. Polonsky, Phys. Rev. D **60**, 115011 (1999); S. Moretti and D. P. Roy, Phys. Lett. B **470**, 209 (1999); D.J. Miller, S. Moretti, D.P. Roy and W.J. Stirling, Phys. Rev. **D61**, 055011 (2000); A. A. Barrientos Bendezu and B. A. Kniehl, Phys. Rev. D **59**, 015009 (1999); Phys. Rev. D **61**, 097701 (2000); Phys. Rev. D **63**, 015009 (2001); O. Brein and W. Hollik, Eur. Phys. J. C **13**, 175 (2000); O. Brein, W. Hollik and S. Kanemura, Phys. Rev. D **63**, 095001 (2001); M. Bisset, M. Guchait and S. Moretti, Eur. Phys. J. C **19**, 143 (2001); Z. Fei *et al.*, Phys. Rev. D **63**, 015002 (2001); A. Belyaev, D. Garcia, J. Guasch and J. Sola, JHEP **0206**, 059 (2002); Phys. Rev. D **65**, 031701 (2002); M. Bisset, F. Moortgat and S. Moretti, Eur. Phys. J. C **30**, 419 (2003); Q. H. Cao, S. Kanemura and C. P. Yuan, Phys. Rev. D **69**, 075008 (2004); E. Asakawa, O. Brein and S. Kanemura, Phys. Rev. D **72**, 055017 (2005); E. Asakawa and S. Kanemura, Phys. Lett. B **626**, 111 (2005).
- [26] A. Abulencia *et al.* [CDF Collaboration], Phys. Rev. Lett. **96**, 042003 (2006).
- [27] For a review see: F. Borzumati and A. Djouadi, Phys. Lett. B **549**, 170 (2002); D. P. Roy, Mod. Phys. Lett. A **19**, 1813 (2004).
- [28] C. Amsler *et al.* [Particle Data Group], Phys. Lett. B **667**, 1 (2008).
- [29] S. Kanemura, T. Kubota and E. Takasugi, Phys. Lett. B **313**, 155 (1993); J. Horejsi and M. Kladiva, Eur. Phys. J. C **46**, 81 (2006); A. G. Akeroyd, A. Arhrib and E. M. Naimi, Phys. Lett. B **490**, 119 (2000).
- [30] H. J. He and C. P. Yuan, Phys. Rev. Lett. **83**, 28 (1999).
- [31] B. Dudley and C. Kolda, arXiv:0901.3337 [hep-ph].
- [32] F. Borzumati and C. Greub, Phys. Rev. D **59**, 057501 (1999).
- [33] Z. J. Xiao and L. Guo, Phys. Rev. D **69**, 014002 (2004).
- [34] N. G. Deshpande, P. Lo, J. Trampetic, G. Eilam and P. Singer, Phys. Rev. Lett. **59**, 183 (1987).
- [35] D. Bowser-Chao, K. m. Cheung and W. Y. Keung, Phys. Rev. D **59**, 115006 (1999) [arXiv:hep-ph/9811235].

- [36] A. Wahab El Kaffas, P. Osland and O. M. Ogreid, Phys. Rev. D **76**, 095001 (2007) [arXiv:0706.2997 [hep-ph]].
- [37] G. Isidori, arXiv:0710.5377 [hep-ph].
- [38] J. L. Diaz-Cruz *et al.*, in preparation.
- [39] F. Mahmoudi and O. Stal, arXiv:0907.1791 [hep-ph].
- [40] P. H. Chankowski, M. Krawczyk and J. Zochowski, Eur. Phys. J. C **11**, 661 (1999) [arXiv:hep-ph/9905436].
- [41] K. Fujikawa, Phys. Rev. D **7**, 393 (1973); M. Bace and N. D. Hari Dass, Ann. Phys. (NY) **94**, 349 (1975); M. B. Gavela, G. Girardi, C. Malleville, and P. Sorba, Nucl. Phys. **B193**, 257 (1981); N. M. Monyonko, J. H. Reid, and A. Sen, Phys. Lett. B **136**, 265 (1984); N. M. Monyonko and J. H. Reid, Phys. Rev. D **32**, 962 (1985); J. M. Hernández, M. A. Pérez, G. Tavares-Velasco, and J. J. Toscano, Phys. Rev. D **60**, 013004 (1999).
- [42] G. Passarino and M. Veltman, Nucl. Phys. **B160**, 151 (1979).
- [43] G. J. van Oldenborgh, J. A. M. Vermaseren, Z. Phys. C **46**, 425 (1990); T. Hahn and M. Pérez-Victoria, Comput. Phys. Commun. **118**, 153 (1999).
- [44] S. Moretti, Pramana **60**, 369 (2003).
- [45] S. Moretti, J. Phys. G **28**, 2567 (2002).